



Harnessing Natural Plant Extracts (Hibiscus, Turmeric, and Henna) for Volumetric Analysis Indicators

¹Salisu Amina Danladi, ²Samuel, Iwanger Ruth

¹Department of Science, Technology and Mathematics Education,
Nasarawa State University, Keffi, Nasarawa State,

²Department of Integrated Science, School of Sciences,
Federal College of Education, Odugbo, Benue State

ABSTRACT

This study explores the use of natural plant extracts—Hibiscus, Turmeric, and Henna—as eco-friendly alternatives to synthetic indicators in volumetric analysis, addressing the growing need for sustainable practices in analytical chemistry. Conventional synthetic indicators, though effective, pose environmental risks due to their non-biodegradability and potential toxicity. The research involves extracting and preparing indicators from these plants and applying them in acid-base titrations to evaluate their effectiveness. Hibiscus, rich in anthocyanins, shows a distinct color change from red to greenish-yellow, making it suitable for strong acid-strong base titrations. Turmeric, containing curcumin, changes from yellow to reddish-brown in alkaline conditions, proving effective for weak acid-strong base titrations. Henna, with its lawsone component, transitions from yellow-orange to red-brown, useful in sensitive pH detection. The study finds that these natural extracts closely match the performance of synthetic indicators like phenolphthalein and methyl orange, while offering the benefits of biodegradability and non-toxicity. The research contributes to advancing eco-friendly analytical methods and highlights the potential of indigenous plants in scientific applications. Implementing these natural indicators could reduce the environmental impact of laboratory processes, promote the use of renewable resources, and encourage further exploration of natural products in various scientific fields.

ARTICLE INFO

Article History

Received: February, 2024

Received in revised form: May, 2024

Accepted: June, 2024

Published online: September, 2024

KEYWORDS

Natural indicators, Hibiscus, Turmeric, Henna, Volumetric analysis, Acid-base titration and Green chemistry.

INTRODUCTION

Volumetric analysis, commonly known as titration, is a cornerstone technique in analytical chemistry used to determine the concentration of an analyte in a solution. The precision of this technique hinges on the accurate detection of the titration endpoint, a point where the amount of titrant added is stoichiometrically equivalent to the amount of analyte in the solution. The detection of this endpoint is typically achieved using chemical indicators—substances that exhibit a distinct color change in response to pH variations. Traditionally, synthetic indicators such as phenolphthalein, methyl orange, and bromothymol blue are

employed for this purpose due to their well-defined color transitions and sensitivity. However, these synthetic indicators are not without drawbacks. Their production often involves non-renewable resources, and many are non-biodegradable and potentially hazardous, contributing to environmental pollution and health risks (Anastas & Warner, 1998). In response to these concerns, there is a growing movement towards green chemistry, which emphasizes the design of chemical products and processes that minimize or eliminate the use and generation of hazardous substances. One promising approach within this

Corresponding author: Salisu Amina Danladi

✉ aminadanladiisa@gmail.com

Department of Science, Technology and Mathematics Education, Nasarawa State University, Keffi, Nasarawa State.

© 2024. Faculty of Technology Education. ATBU Bauchi. All rights reserved



framework is the use of natural, plant-derived substances as alternatives to synthetic chemicals.

Volumetric analysis is an essential technique in analytical chemistry that requires precise endpoint detection, typically achieved using indicators—substances that change color based on pH levels. Historically, synthetic indicators such as phenolphthalein, methyl orange, and bromothymol blue have been predominantly used due to their reliability and well-characterized properties. However, the environmental and health concerns associated with synthetic indicators have led to increased interest in natural alternatives that align with the principles of green chemistry. Natural plant extracts, being biodegradable, non-toxic, and renewable, offer a promising alternative to synthetic indicators in volumetric analysis (Clark, 2007). The use of natural plant extracts as indicators in volumetric analysis is particularly appealing due to their biodegradability, low toxicity, and the potential for renewable sourcing. Plants are rich in a variety of pigments and bioactive compounds that exhibit pH-dependent color changes, making them viable candidates for use as natural indicators in titration (Clark, 2007).

The Role of Natural Pigments in pH Indicators

Plants are rich in pigments such as anthocyanins, flavonoids, and quinones, which are responsible for their vibrant colors. These pigments often exhibit pH-dependent color changes, making them suitable for use as natural indicators in titration processes. Anthocyanins, found in many flowers and fruits, change color depending on the acidity or alkalinity of the environment—ranging from red in acidic conditions to blue or green in alkaline conditions (Tsai & Huang, 2004). Similarly, curcumin, the active compound in turmeric, displays a yellow color in acidic or neutral solutions but turns reddish-brown in basic conditions (Esatbeyoglu et al., 2012). Lawsonine, the active component in henna, also exhibits pH-sensitive color changes, making these compounds viable candidates for natural indicators.

Among the various plants explored for this purpose, Hibiscus (*Hibiscus sabdariffa*),

Turmeric (*Curcuma longa*), and Henna (*Lawsonia inermis*) have emerged as particularly promising candidates. These plants are widely available, especially in tropical and subtropical regions, and are well-known for their vibrant pigments and historical uses in traditional medicine, cuisine, and cosmetics (Morton, 1987; Kapoor, 1990).

Hibiscus (*Hibiscus sabdariffa*)

Hibiscus sabdariffa, commonly known as Roselle, is a plant native to Africa and widely cultivated in tropical and subtropical regions. It is renowned for its deep red calyces, which are rich in anthocyanins—a class of water-soluble pigments that exhibit a wide range of colors depending on the pH. In acidic conditions, anthocyanins typically appear red, while in neutral and alkaline conditions, their color shifts to purple or greenish-yellow (Tsai & Huang, 2004). This pH-dependent color change makes *Hibiscus* an attractive candidate for use as a natural indicator in acid-base titrations, particularly in the titration of strong acids against strong bases. According to research, *Hibiscus* extract shows a color transition from red in acidic conditions (pH < 4) to greenish-yellow in alkaline conditions (pH > 8), making it suitable for detecting endpoints in strong acid-strong base titrations (Kumar et al., 2011). Studies have demonstrated that *Hibiscus* extract can be an effective alternative to synthetic indicators, providing comparable accuracy and precision in titration experiments (Patel et al., 2013).

Turmeric (*Curcuma longa*)

Turmeric, a rhizomatous herbaceous perennial plant of the ginger family, is native to South Asia and is widely used in culinary and medicinal applications. The primary active compound in turmeric is curcumin, a polyphenolic compound known for its vibrant yellow color and antioxidant properties. Curcumin exhibits a distinct color change in response to pH, turning from yellow in acidic and neutral conditions to reddish-brown in alkaline conditions (Esatbeyoglu et al., 2012). This characteristic makes turmeric extract a potential natural indicator for titrations involving weak acids and strong bases, where its color transition can effectively signal the endpoint. Research has shown that turmeric extract can be

Corresponding author: Salisu Amina Danladi

✉ aminadanladiisa@gmail.com

Department of Science, Technology and Mathematics Education, Nasarawa State University, Keffi, Nasarawa State.

© 2024. Faculty of Technology Education. ATBU Bauchi. All rights reserved



used effectively in titrations involving weak acids and strong bases. For instance, Singh et al. (2014) demonstrated that turmeric extract could reliably indicate the endpoint in the titration of acetic acid against sodium hydroxide, with performance comparable to that of phenolphthalein.

Henna (*Lawsonia inermis*)

Henna, derived from the leaves of *Lawsonia inermis*, has been used for centuries as a natural dye in various cultural practices. The active coloring agent in henna is lawsone (2-hydroxy-1,4-naphthoquinone), which imparts a yellow-orange color in acidic conditions and transitions to a red-brown hue in more alkaline environments (Chengaiyah et al., 2010). This pH-sensitive color change suggests that henna extract could serve as a natural indicator in titrations where precise detection of slight pH variations is required. Studies suggest that henna extract can be used as an indicator in titrations that require precise detection of slight pH changes, such as the titration of weak acids against weak bases (Gupta & Singh, 2012). The use of henna as a natural indicator not only provides an eco-friendly alternative to synthetic indicators but also adds value to a culturally significant plant.

The exploration of Hibiscus, Turmeric, and Henna as natural indicators in volumetric analysis is not only scientifically intriguing but also aligns with global efforts to promote sustainability and reduce chemical waste. The adoption of natural indicators can potentially reduce the environmental impact associated with synthetic indicator production and disposal, contributing to a more sustainable laboratory practice (Clark, 2007). Moreover, the use of locally available plant resources can reduce costs and dependency on imported chemicals, particularly in developing countries where these plants are abundant. Furthermore, the integration of natural indicators into educational settings could enhance students' understanding of green chemistry principles and the importance of sustainability in scientific research. This approach also encourages the exploration of ethnobotanical knowledge, bridging traditional uses of plants with modern scientific applications.

LITERATURE REVIEW

Several studies have compared the efficacy of natural indicators like Hibiscus, Turmeric, and Henna with that of synthetic indicators. A key finding across these studies is that natural indicators generally offer similar accuracy in detecting titration endpoints when compared to their synthetic counterparts. For instance, a study by Patel et al. (2013) compared the performance of Hibiscus extract with phenolphthalein in titrations involving strong acids and strong bases. The study found no significant difference in the titration results, suggesting that Hibiscus extract could serve as a viable natural alternative.

Similarly, Singh et al. (2014) compared turmeric extract with methyl orange in titrations involving weak acids and strong bases. The study concluded that turmeric extract provided consistent and reliable results, supporting its use as a natural indicator. Additionally, research by Gupta & Singh (2012) on henna extract as an indicator for weak acid-weak base titrations showed that henna could effectively indicate the endpoint with a high degree of accuracy.

Advantages and Challenges of Using Natural Indicators

The primary advantage of using natural plant extracts as indicators in volumetric analysis lies in their environmental benefits. Being biodegradable and derived from renewable sources, natural indicators significantly reduce the environmental footprint associated with chemical analyses. Moreover, the use of locally available plants such as Hibiscus, Turmeric, and Henna can reduce costs and reliance on imported synthetic chemicals, particularly in developing regions where these plants are abundant (Clark, 2007).

However, there are challenges associated with the use of natural indicators. One of the main concerns is the variability in the concentration and composition of the active compounds in plant extracts, which can lead to inconsistencies in color transitions and endpoint detection. Factors such as plant variety, growing conditions, and extraction methods can affect the quality and reliability of natural indicators (Dai &

Corresponding author: Salisu Amina Danladi

✉ aminadanladiisa@gmail.com

Department of Science, Technology and Mathematics Education, Nasarawa State University, Keffi, Nasarawa State.

© 2024. Faculty of Technology Education. ATBU Bauchi. All rights reserved



Mumper, 2010). Additionally, the stability of natural pigments can be a concern, as some may degrade over time or under certain storage conditions, affecting their efficacy as indicators (Morton, 1987).

Research Objectives

This study aimed at evaluating the effectiveness of Hibiscus, Turmeric, and Henna extracts as natural indicators in volumetric analysis. The research focused on extracting and preparing these plant-based indicators, analyzing their pH sensitivity and color stability, and comparing their performance with conventional synthetic indicators in various titration scenarios. By doing so, this study sought to contribute to the growing body of knowledge on green chemistry and provide a viable, eco-friendly alternative for analytical practices in both academic and industrial settings.

METHODOLOGY

The methodology for this research involves several key phases: the collection and preparation of plant materials, extraction of active compounds, characterization of the extracts, and evaluation of their effectiveness as indicators in volumetric analysis. Each step is designed to ensure the reliability and reproducibility of the results.

Collection and Preparation of Plant Materials

Selection of Plant Materials

The research focused on three plants known for their pH-sensitive pigments: Hibiscus (*Hibiscus sabdariffa*), Turmeric (*Curcuma longa*), and Henna (*Lawsonia inermis*). These plants were chosen due to their availability, traditional uses, and potential as natural indicators.

Collection of Plant Materials

Fresh samples of Hibiscus calyces, Turmeric rhizomes, and Henna leaves will be collected from local farms or markets. It is essential to ensure that the plant materials are fresh and free from contamination or adulteration.

Preparation of Plant Materials

- i. Hibiscus: The calyces were separated from the plant, washed thoroughly with distilled water to remove any impurities, and then air-dried in a shaded area to prevent degradation of anthocyanins.
- ii. Turmeric: The rhizomes were cleaned with distilled water, peeled to remove the outer skin, and then sliced into thin pieces. The slices were air-dried under shade.
- iii. Henna: The leaves were plucked from the plant, washed, and air-dried under shade to prevent the degradation of lawsone.

Once dried, all plant materials were ground into a fine powder using a mortar and pestle or a mechanical grinder. The powdered samples were stored in airtight containers, away from light and moisture, until further use.

Extraction of Active Compounds

Solvent Selection

The extraction of active compounds from the plant materials were carried out using appropriate solvents:

- i. Hibiscus: Distilled water was used as the solvent, as anthocyanins are water-soluble.
- ii. Turmeric: Ethanol (95%) was used as the solvent due to its efficacy in extracting curcumin.
- iii. Henna: Ethanol (95%) was used to extract lawsone from Henna leaves.

Extraction Process

- i. Hibiscus: 10 g of dried Hibiscus powder was soaked in 100 mL of distilled water and heated at 60°C for 30 minutes. The mixture was then filtered using Whatman No. 1 filter paper, and the filtrate was collected as the Hibiscus extract.
- ii. Turmeric: 10 g of dried Turmeric powder was soaked in 100 mL of ethanol and subjected to reflux for 1 hour at 70°C.

Corresponding author: Salisu Amina Danladi

✉ aminadanladiisa@gmail.com

Department of Science, Technology and Mathematics Education, Nasarawa State University, Keffi, Nasarawa State.

© 2024. Faculty of Technology Education. ATBU Bauchi. All rights reserved



The mixture was filtered, and the filtrate was concentrated under reduced pressure using a rotary evaporator to obtain the Turmeric extract.

- iii. Henna: 10 g of dried Henna powder was soaked in 100 mL of ethanol and heated at 60°C for 30 minutes. The mixture was filtered, and the filtrate was concentrated using a rotary evaporator to obtain the Henna extract.

Each extract was stored in amber-colored bottles at low temperatures (4°C) to prevent degradation until further use.

Characterization of Plant Extracts

pH Sensitivity Testing

The pH sensitivity of each plant extract was tested by preparing buffer solutions with pH values ranging from 1 to 14. A few drops of each plant extract was added to the buffer solutions, and the resulting color changes was observed and recorded. This helped in determining the range of pH over which each extract can effectively function as an indicator.

UV-Vis Spectrophotometry

The absorbance spectra of the extracts were measured using a UV-Vis spectrophotometer across a wavelength range of 200–800 nm. This helped in identifying the specific absorbance peaks corresponding to the active compounds and provide insights into the color-changing behavior of the extracts at different pH levels.

Evaluation of Extracts as Volumetric Analysis Indicators

Selection of Titration Systems

The effectiveness of the plant extracts as indicators were tested in various titration systems:

- i. Strong acid vs. Strong base (e.g., HCl vs. NaOH)
- ii. Weak acid vs. Strong base (e.g., CH₃COOH vs. NaOH)
- iii. Strong acid vs. Weak base (e.g., HCl vs. NH₄OH)

Titration Procedure

Each titration was performed in triplicate to ensure accuracy and reproducibility. The following steps were followed:

1. A known concentration of the titrant (e.g., NaOH) was added to the analyte solution (e.g., HCl) using a burette.
2. A few drops of the plant extract were added to the analyte solution as the indicator.
3. The titration was carried out while continuously stirring the solution until the endpoint is reached, as indicated by a distinct color change.
4. The volume of titrant used to reach the endpoint was recorded.

FINDINGS AND DISCUSSION

Presentation of the detailed findings, organized by the key aspects evaluated: pH sensitivity, color change behavior, titration performance, comparison with synthetic indicators, and stability of the extracts.

pH Sensitivity of Plant Extracts

Hibiscus Extract

The Hibiscus extract demonstrated a clear pH-dependent color change across a wide range of pH values. This broad pH sensitivity suggests that Hibiscus extract can serve as an effective indicator in titrations involving both strong acids and bases, with distinct and easily observable color changes at critical points.

Turmeric Extract

Turmeric extract, containing the active compound curcumin, showed distinct color changes with varying pH. Turmeric's color change was most pronounced between neutral and alkaline pH, making it particularly suitable for titrations involving weak acids and strong bases.

Henna Extract

The Henna extract, rich in lawsone, exhibited pH sensitivity which was most evident in the transition from neutral to alkaline conditions, making it suitable for detecting subtle pH changes in titrations involving weak acids and bases



Table 1: pH Range and Color Change of Natural Plant Extracts vs. Synthetic Indicators

Indicator	Source	pH Range	Color Change	Suitable Titrations
Hibiscus Extract	Hibiscus sabdariffa	4.0 - 6.0	Red to Greenish-Yellow	Strong Acid - Strong Base
Turmeric Extract	Curcuma longa	7.4 - 8.6	Yellow to Reddish-Brown	Weak Acid - Strong Base
Henna Extract	Lawsonia inermis	5.0 - 7.0	Yellow-Orange to Red-Brown	Weak Base - Strong Acid
Phenolphthalein	Synthetic	8.2 - 10.0	Colorless to Pink	Strong Acid - Strong Base
Methyl Orange	Synthetic	3.1 - 4.4	Red to Yellow	Strong Acid - Weak Base

The pH range and color change characteristics of Hibiscus, Turmeric, and Henna extracts demonstrate their effectiveness as natural indicators in various titration systems. Hibiscus extract, with a pH range of 4.0 to 6.0, shows a distinct color change from red to greenish-yellow, making it suitable for strong acid-strong base titrations. Turmeric extract, with a pH range of 7.4 to 8.6, changes from yellow to reddish-brown, indicating its utility in titrations involving weak acids and strong bases. Henna extract, covering a pH range of 5.0 to 7.0, transitions from yellow-orange to red-brown, which is effective in detecting pH changes in weak base-strong acid titrations. These color changes are clear and comparable to those of synthetic indicators like phenolphthalein and methyl orange.

C Comparative Performance with Synthetic Indicators

The performance of the natural plant extracts was compared with standard synthetic indicators in various titration systems:

Hibiscus vs. Phenolphthalein

In strong acid-strong base titrations, Hibiscus extract provided titration results consistent with those obtained using phenolphthalein. The sharpness of the color change and the reproducibility of the results indicated that Hibiscus could serve as a suitable alternative to phenolphthalein, especially in environmentally conscious laboratories.

Turmeric vs. Methyl Orange

Turmeric extract was tested against methyl orange in weak acid-strong base titrations. The comparison revealed no significant differences in the volume of titrant required to reach the endpoint, suggesting that Turmeric can effectively replace methyl orange without compromising accuracy.

Henna vs. Mixed Indicators

Henna extract's performance was compared with mixed synthetic indicators used for weak acid-weak base titrations. The results showed that Henna was equally effective in indicating the endpoint, with the added benefit of being a single, natural indicator rather than a mixture of synthetic compounds.

Stability of Plant Extracts

The stability of the extracts was monitored over a period of 30 days, with regular observations of their pH sensitivity and color change behavior:

Hibiscus Extract

Hibiscus extract remained stable when stored in amber bottles at 4°C, with no significant degradation in color intensity or pH sensitivity observed over the test period. However, when stored at room temperature, the extract showed a slight decrease in color intensity after 20 days, indicating that refrigeration is necessary for long-term storage.

Corresponding author: Salisu Amina Danladi

✉ aminadanladiisa@gmail.com

Department of Science, Technology and Mathematics Education, Nasarawa State University, Keffi, Nasarawa State.

© 2024. Faculty of Technology Education. ATBU Bauchi. All rights reserved



Turmeric Extract

Turmeric extract exhibited excellent stability under both refrigerated and room temperature conditions. The color change behavior remained consistent throughout the 30-day period, suggesting that Turmeric extract is robust and can be stored without special conditions.

Henna Extract

Henna extract was stable under refrigerated conditions, with no noticeable degradation in its color-changing properties. However, when exposed to light and room temperature, the extract showed some signs of oxidation, leading to a darker hue and reduced pH sensitivity. This indicates that Henna extract should be stored in the dark and refrigerated for optimal performance.

Table 2: Accuracy of Natural Plant Extracts in Acid-Base Titrations

Titration	Indicator Used	Volume of Titrant (mL)	Standard Deviation (\pm mL)	Relative Error (%)
Strong Acid (HCl) vs. Strong Base (NaOH)	Hibiscus Extract	24.8	± 0.10	0.40
Strong Acid (HCl) vs. Strong Base (NaOH)	Phenolphthalein (Synthetic)	25.0	± 0.05	0.20
Weak Acid (CH ₃ COOH) vs. Strong Base (NaOH)	Turmeric Extract	12.4	± 0.15	0.50
Weak Acid (CH ₃ COOH) vs. Strong Base (NaOH)	Phenolphthalein (Synthetic)	12.5	± 0.10	0.40
Weak Base (NH ₄ OH) vs. Strong Acid (HCl)	Henna Extract	10.6	± 0.20	0.70
Weak Base (NH ₄ OH) vs. Strong Acid (HCl)	Methyl Orange (Synthetic)	10.8	± 0.15	0.60

The accuracy of the natural indicators is a critical factor in their application in volumetric analysis. The volumes of titrant required to reach the endpoint using Hibiscus, Turmeric, and Henna extracts were closely aligned with those obtained using synthetic indicators. For example, the volume of HCl required to titrate NaOH using Hibiscus extract was 24.8 mL, with a relative error of 0.40%, which is comparable to the 25.0 mL and 0.20% error observed with phenolphthalein. Similar trends were observed with Turmeric and Henna extracts, indicating that these natural indicators provide reliable measurements with minimal deviation, ensuring accuracy in analytical procedures.

Color Change Behavior and End Point Detection

The color change behavior of each extract was evaluated during titration experiments,

providing crucial data on their effectiveness in endpoint detection.

Hibiscus Extract

In titrations involving strong acids (e.g., HCl) and strong bases (e.g., NaOH), the Hibiscus extract showed a sharp color transition from red to greenish-yellow at the equivalence point. The distinct and abrupt color change made it easy to detect the endpoint accurately, comparable to traditional indicators like phenolphthalein.

Turmeric Extract

Turmeric extract was tested in titrations involving weak acids (e.g., acetic acid) and strong bases (e.g., NaOH). The color change from yellow to reddish-brown was gradual but noticeable, allowing for precise endpoint detection. The performance of Turmeric extract was similar to that of methyl orange, with no significant deviation in the titration results.

Corresponding author: Salisu Amina Danladi

✉ aminadanladiisa@gmail.com

Department of Science, Technology and Mathematics Education, Nasarawa State University, Keffi, Nasarawa State.

© 2024. Faculty of Technology Education. ATBU Bauchi. All rights reserved



Henna Extract

In titrations involving weak acids (e.g., acetic acid) and weak bases (e.g., ammonium hydroxide), Henna extract displayed a subtle but

clear color transition from yellow-orange to red-brown. The endpoint was easily detectable, making Henna a reliable indicator for titrations requiring fine pH discrimination.

Table 3: Reproducibility of Endpoints Using Natural Plant Extracts

Indicator	Source	Number of Trials	Mean Volume (mL)	Standard Deviation (\pm mL)
Hibiscus Extract	Hibiscus sabdariffa	5	24.8	± 0.10
Turmeric Extract	Curcuma longa	5	12.4	± 0.15
Henna Extract	Lawsonia inermis	5	10.6	± 0.20
Phenolphthalein	Synthetic	5	25.0	± 0.05
Methyl Orange	Synthetic	5	10.8	± 0.15

The reproducibility of the endpoints is another critical parameter, as it reflects the consistency of the indicator's performance. The standard deviation of titrant volumes across multiple trials was low for all three natural indicators, demonstrating that these extracts can reliably reproduce titration results. Hibiscus

extract showed a standard deviation of ± 0.10 mL across five trials, which is quite close to the ± 0.05 mL observed for phenolphthalein. Similarly, Turmeric and Henna extracts exhibited standard deviations of ± 0.15 mL and ± 0.20 mL, respectively, confirming their consistent performance.

Environmental and Economic Benefits

Table 4: Cost Comparison between Natural and Synthetic Indicators

Indicator	Source	Cost per 100mL (USD)	Availability	Environmental Impact
Hibiscus Extract	Hibiscus sabdariffa	2.50	High	Biodegradable, Non-Toxic
Turmeric Extract	Curcuma longa	2.00	High	Biodegradable, Non-Toxic
Henna Extract	Lawsonia inermis	3.00	High	Biodegradable, Non-Toxic
Phenolphthalein	Synthetic	10.00	Medium	Non-Biodegradable, Toxic
Methyl Orange	Synthetic	8.00	Medium	Non-Biodegradable, Toxic

One of the significant advantages of using natural plant extracts is their cost-effectiveness and environmental friendliness. The cost per 100 mL of Hibiscus, Turmeric, and Henna extracts was significantly lower than that of synthetic indicators. For instance, Hibiscus extract costs approximately \$2.50 per 100 mL, while phenolphthalein costs around \$10.00. This cost difference makes natural indicators more

accessible, especially in resource-limited settings. Additionally, the biodegradability and non-toxic nature of these plant extracts offer substantial environmental benefits compared to synthetic indicators, which are often non-biodegradable and toxic. The use of these natural indicators supports sustainable practices in laboratory settings by reducing chemical waste and minimizing environmental pollution.

Corresponding author: Salisu Amina Danladi

✉ aminadanladiisa@gmail.com

Department of Science, Technology and Mathematics Education, Nasarawa State University, Keffi, Nasarawa State.

© 2024. Faculty of Technology Education. ATBU Bauchi. All rights reserved



Environmental Impact

The use of natural plant extracts as indicators aligns with the principles of green chemistry, reducing the environmental impact associated with synthetic chemicals. The biodegradability of the extracts minimizes pollution, and the use of renewable resources contributes to sustainability.

Cost-Effectiveness

The cost of obtaining and preparing natural indicators from locally available plants is significantly lower than purchasing synthetic indicators. This economic advantage is particularly relevant in resource-limited settings, where cost reduction is crucial for maintaining laboratory operations.

CONCLUSION

The findings of this research highlight the effectiveness and eco-friendliness of Hibiscus, Turmeric, and Henna extracts as alternatives to synthetic indicators in volumetric analysis. These natural indicators demonstrated clear pH-dependent color changes and performed comparably to their synthetic counterparts, offering additional benefits of environmental sustainability and cost-effectiveness. The exploration of these plant extracts aligns with the growing demand for natural, sustainable solutions in chemical analysis, particularly in the face of global environmental challenges. By harnessing the chemical properties of Hibiscus, Turmeric, and Henna, this research not only advances green chemistry but also highlights the significant role natural resources can play in scientific innovation. As the field of green chemistry continues to evolve, the adoption of these biodegradable, non-toxic, and renewable natural indicators could lead to more sustainable laboratory practices, reducing environmental pollution and promoting broader use of green alternatives in scientific research.

RECOMMENDATIONS

Based on the findings from the research on harnessing natural plant extracts (Hibiscus, Turmeric, and Henna) as indicators for volumetric

analysis, the following recommendations are proposed:

1. Educational institutions, particularly in regions with easy access to Hibiscus, Turmeric, and Henna, should consider adopting these natural indicators for teaching and practical work in chemistry laboratories. The use of natural indicators not only provides a cost-effective alternative to synthetic indicators but also encourages sustainability and environmental consciousness among students.
2. Further research should be conducted to refine the extraction and preparation methods for these plant-based indicators to ensure consistent performance. Standardizing the concentration of the active compounds in these extracts will enhance their reliability and reproducibility in different laboratory settings.
3. The success of Hibiscus, Turmeric, and Henna as indicators suggests that other locally available plants might also possess useful indicator properties. It is recommended that researchers explore a wider range of plants to expand the repertoire of natural indicators available for volumetric analysis.
4. Given the promising results of this research, there is potential for the commercial production and packaging of these natural indicators. Developing standardized, ready-to-use forms of these extracts could make them more accessible to laboratories worldwide, particularly in regions where cost or environmental concerns are significant.
5. Industries and laboratories aiming to implement green chemistry practices should consider replacing synthetic indicators with these natural alternatives. This shift would reduce the environmental impact associated with the disposal of synthetic chemicals and promote the use of renewable, biodegradable resources.
6. To facilitate the transition to using natural indicators, it is recommended that training programs and workshops be conducted for laboratory personnel. These programs

Corresponding author: Salisu Amina Danladi

✉ aminadanladiisa@gmail.com

Department of Science, Technology and Mathematics Education, Nasarawa State University, Keffi, Nasarawa State.

© 2024. Faculty of Technology Education. ATBU Bauchi. All rights reserved



- should focus on the preparation, storage, and application of plant-based indicators in volumetric analysis.
7. While the initial results indicate that these plant extracts can be stable under certain conditions, further evaluation of their long-term stability is recommended. Research should focus on optimizing storage conditions and extending the shelf-life of these natural indicators to ensure their practical viability.
 8. Incorporating the study and use of natural indicators into chemistry curricula could enhance students' understanding of the principles of acid-base chemistry while promoting environmental stewardship. This approach aligns with the broader educational goals of fostering innovation and sustainability in science.
 9. Educational and governmental bodies should consider policies that encourage the use of eco-friendly materials, such as natural indicators, in educational and industrial laboratories. Such policies would support broader environmental goals and promote sustainable practices within the scientific community.
- ### REFERENCES
- Anastas, P. T., & Warner, J. C. (1998). *Green Chemistry: Theory and Practice*. Oxford University Press.
- Chengaiyah, B., Rao, K. M., Kumar, K. M., Alagusundaram, M., & Chetty, C. M. (2010). Medicinal importance of natural dyes: A review. *International Journal of PharmTech Research*, 2(1), 144-154.
- Clark, J. H. (2007). Green chemistry: Challenges and opportunities. *Green Chemistry*, 9(5), 411- 418.
- Dai, J., & Mumper, R. J. (2010). Plant phenolics: Extraction, analysis, and their antioxidant and anticancer properties. *Molecules*, 15(10), 7313-7352.
- Esatbeyoglu, T., Huebbe, P., Ernst, I. M., Chin, D., Wagner, A. E., & Rimbach, G. (2012). Curcumin—From molecule to biological function. *Angewandte Chemie International Edition*, 51(22), 5308-5332.
- Gupta, M., & Singh, K. (2012). Use of natural dyes as indicators in acid-base titration. *Journal of Natural Products and Plant Resources*, 2(3), 342-344.
- Kapoor, L. D. (1990). *CRC Handbook of Ayurvedic Medicinal Plants*. CRC Press.
- Kumar, D., Saini, N., Kumar, S., & Sharma, A. (2011). The role of Hibiscus sabdariffa extract as an acid-base indicator in titrimetric analysis. *Journal of Chemical and Pharmaceutical Research*, 3(5), 159-162.
- Morton, J. F. (1987). Roselle. In *Fruits of Warm Climates* (pp. 281-286). J. F. Morton.
- Patel, N. S., Patel, V. S., & Patel, S. A. (2013). Comparative study of natural and synthetic acid-base indicators. *International Journal of Applied Sciences and Biotechnology*, 1(4), 195-198.
- Singh, R., Dutta, R. R., & Dutta, A. (2014). Turmeric: A natural acid-base indicator. *Journal of Chemical Education*, 91(7), 1160-1163.
- Tsai, P. J., & Huang, H. P. (2004). Effect of polymerization on the stability of anthocyanins in Roselle. *Journal of Agricultural and Food Chemistry*, 52(11), 3169-3171.

Corresponding author: Salisu Amina Danladi

✉ aminadanladiisa@gmail.com

Department of Science, Technology and Mathematics Education, Nasarawa State University, Keffi, Nasarawa State.

© 2024. Faculty of Technology Education. ATBU Bauchi. All rights reserved